## **United States** Health Department Website: hhs.gov

### Summary

### **Highlights**


#### **Drug Deaths**

from 6.9 to 9.9 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

#### **Number of Geriatric Providers**

from 31.4 to 36.4 per 100,000 adults ages 65+ between September 2019 and September 2022.

#### **Multiple Chronic Conditions**



from 46% to 52% of Medicare beneficiaries ages 65+ between 2020 and 2021.

#### **Poverty**

from 9.4% to 10.3% of adults ages 65+ between 2019 and 2021.

#### **High-speed Internet**



from 78.0% to 83.1% of households with adults ages 65+ between 2019 and 2021.

**Food Insecurity** 



from 12.6% to 11.9% of adults ages 60+ between 2019 and 2020.

### Measures

leasure	S	Value
Social & Econo		
Community and		13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	11.8
Economic Resources	Food Insecurity (% of adults ages 60+)	11.9%
Resources	Poverty (% of adults ages 65+)	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	81.0
Social Support and	Community Support Expenditures (dollars per adult ages 60+)	\$62
Engagement	High-speed Internet (% of households with adults ages 65+)	83.1%
	Low-care Nursing Home Residents (% of residents)	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	_
	Volunteerism (% of adults ages 65+)	22.1%
Physical Enviro	onment*	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	7.8
Quality	Drinking Water Violations (% of community water systems)	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	32.3%
Clinical Care*		
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	67.5%
bervices	Pneumonia Vaccination (% of adults ages 65+)	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	96.1%
Care	Hospice Care (% of Medicare decedents)	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	1,482
Behaviors*		
Nutrition	Exercise (% of adults ages 65+)	23.1%
and Physical Activity	Fruit and Vegetable Consumption (% of adults ages 65+)	7.3%
ACTIVITY	Physical Inactivity (% of adults ages 65+ in fair or better health)	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	26.0%
lobacco Use	Smoking (% of adults ages 65+)	8.9%
Health Outcon	nes*	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	9.9
	Excessive Drinking (% of adults ages 65+)	7.0%
	Frequent Mental Distress (% of adults ages 65+)	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	2,151
	Early Death Racial Disparity (ratio)***	1.5
Physical Health	Falls (% of adults ages 65+)	27.1%
	Frequent Physical Distress (% of adults ages 65+)	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	52%
	Obesity (% of adults ages 65+)	29.5%

\* Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia. \*\* Disparity measures compare the group with the highest or lowest rate to the white rate.

\* Non-ranking measure. - Data not available, missing or suppressed.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



U.S.

# Hawaii State Health Department Website: health.hawaii.gov

### Summary

### **Strengths**

- Low early death rate
- Low prevalence of obesity
- Low prevalence of physical inactivity

### Challenges

- Low volunteerism rate
- Low home health care worker rate
- High prevalence of severe housing problems

### Highlights

**Home Health Care Workers** 

29%

from 32.8 to 23.4 aides per 1,000 adults ages 65+ between 2020 and 2021.

Food Insecurity



from 10.2% to 7.5% of adults ages 60+ between 2017-2018 and 2019-2020.

#### **Multiple Chronic Conditions**



from 41% to 47% of Medicare beneficiaries ages 65+ between 2020 and 2021.

## Measures

/leasure	15	Rank	Value	Valu
Social & Econo	omic Factors*	7	0.633	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	5	4.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	1	5.8	11.8
Economic Resources	Food Insecurity (% of adults ages 60+)	5	7.5%	11.9%
	Poverty (% of adults ages 65+)	22	9.4%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support and	Community Support Expenditures (dollars per adult ages 60+)	16	\$54	\$62
	High-speed Internet (% of households with adults ages 65+)	9	86.1%	83.1%
ngagement	Low-care Nursing Home Residents (% of residents)	1	3.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	14	42	_
	Volunteerism (% of adults ages 65+)	41	18.4%	22.19
hysical Enviro	onment*	12	0.503	
ir and Water	Air Pollution (micrograms of fine particles per cubic meter)	1	3.8	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)*	41	33.7%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	44	37.0%	32.3
linical Care*		15	0.322	
ccess to Care	Avoided Care Due to Cost (% of adults ages 65+)	26	3.0%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	8	46.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	46	23.4	60.3
reventive	Cancer Screenings (% of adults ages 65-75)	5	80.4%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	35	65.6%	67.5
ervices	Pneumonia Vaccination (% of adults ages 65+)	49	60.4%	69.79
Quality of Care	Dedicated Health Care Provider (% of adults ages 65+)	4	97.4%	96.19
	Hospice Care (% of Medicare decedents)	22	48.0%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	1	73.3%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	1	736	1,482
Behaviors*		25	0.042	
lutrition	Exercise (% of adults ages 65+)	15	25.8%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	28	6.4%	7.3%
	Physical Inactivity (% of adults ages 65+ in fair or better health)	6	25.4%	31.25
ctivity			25.4% 38.6%	
leep Health	better health)	6		26.0
leep Health obacco Use	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	6 50	38.6%	26.0
leep Health obacco Use <b>lealth Outcon</b> tehavioral	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	6 50 4	38.6% 7.0%	26.0° 8.9%
leep Health obacco Use <b>lealth Outcon</b> ehavioral	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) <b>nes*</b>	6 50 4 <b>1</b>	38.6% 7.0% <b>1.132</b>	26.0 8.9%
leep Health obacco Use <b>lealth Outcon</b> ehavioral	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+)	6 50 4 <b>1</b> 40	38.6% 7.0% <b>1.132</b> 9.0%	26.0° 8.9% 7.8% 9.9
leep Health obacco Use <b>lealth Outcon</b> ehavioral	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	6 50 4 <b>1</b> 40 46	38.6% 7.0% <b>1.132</b> 9.0% 15.3	26.0° 8.9% 7.8% 9.9 7.0%
ctivitý leep Health obacco Use <b>lealth Outcon</b> ehavioral lealth	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+)	6 50 4 <b>1</b> 40 46 34	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8%	26.0 <sup>4</sup> 8.9% 7.8% 9.9 7.0% 8.5%
ctivity leep Health obacco Use <b>lealth Outcon</b> ehavioral lealth	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	6 50 4 <b>1</b> 40 46 34 34 10	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1%	26.01 8.9% 7.8% 9.9 7.0% 8.5% 16.9
ctivity leep Health obacco Use <b>lealth Outcon</b> ehavioral lealth fortality	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)** <sup>‡</sup>	6 50 4 <b>1</b> 40 46 34 10 9	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1% 13.8	26.09 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2,15
ctivity leep Health obacco Use <b>lealth Outcon</b> ehavioral lealth fortality	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death (ages 65+) Farly Context (ratio)*** Falls (% of adults ages 65+)	6 50 4 <b>1</b> 40 46 34 10 9 1	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1% 13.8 1.552	26.09 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2,15 1.5
activity ileep Health obacco Use <b>Health Outcon</b> Iehavioral Iealth fortality	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+)	6 50 4 <b>1</b> 40 46 34 10 9 1 1 -	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1% 13.8 1,552 1.5	26.09 8.9% 9.9 7.0% 8.5% 16.9 2,15 1.5 27.1%
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality Physical Health	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death (ages 65+) Farly Context (ratio)*** Falls (% of adults ages 65+)	6 50 4 1 40 46 34 10 9 1 - 2	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1% 13.8 1.552 1.5 21.3%	31.29 26.09 8.9% 9.9 7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8% 52%
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality	better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) <sup>‡</sup> Drug Deaths (deaths per 100,000 adults ages 65+) <sup>‡</sup> Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death (adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	6 50 4 4 40 46 34 10 9 1 - 2 3	38.6% 7.0% <b>1.132</b> 9.0% 15.3 7.8% 7.1% 13.8 1.552 1.5 21.3% 11.9%	26.09 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8%

\* Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

\*\* Disparity measures compare the group with the highest or lowest rate to the white rate.

\* Non-ranking measure.

Data not available, missing or suppressed.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



State State

U.S.